

Extensions of the Scherck-Kemperman Theorem

Y. O. Hamidoune*

September 30, 2009

Abstract

Let $\Gamma = (V, E)$ be a reflexive relation with a transitive automorphisms group. Let $v \in V$ and let F be a finite subset of V with $v \in F$.

We prove that the size of $\Gamma(F)$ (the image of F) is at least

$$|F| + |\Gamma(v)| - |\Gamma^-(v) \cap F|.$$

Let A, B be finite subsets of a group G . Applied to Cayley graphs, our result reduces to following extension of the Scherck-Kemperman Theorem, proved by Kemperman:

$$|AB| \geq |A| + |B| - |A \cap (cB^{-1})|,$$

for every $c \in AB$.

1 Introduction

Let A, B be finite subsets of a group G . The *Minkowsky product* of A by B is defined as $AB = \{xy : x \in A \text{ and } y \in B\}$.

Solving a problem of Moser, Scherck proved in [13] that

$$|AB| \geq |A| + |B| - 1,$$

if G is abelian and if $A \cap B^{-1} = \{1\}$. The validity of this result in the non-abelian case was proved by Kemperman [9]. Kemperman mentioned in [9] that this generalization was independently obtained by Wehn. This result is known as the Scherck-Kemperman Theorem.

*UPMC Univ Paris 06, E. Combinatoire, Case 189, 4 Place Jussieu, 75005 Paris, France.

Kemperman's Theorem [9] states that

$$|AB| \geq |A| + |B| - |A \cap (cB^{-1})|,$$

for every $c \in AB$.

The reader may find several applications of the Scherk-Kemperman Theorem to the Theory of Non-unique factorization in the text book of Geroldinger-Halter-Koch [3]. Recall that this result is used, among other tools, by Olson in [12], to prove that for any subset S of a finite group G with $|S| \geq 3\sqrt{|G|}$, there exist distinct elements $x_1, \dots, x_k \in S$ with $x_1 \cdots x_k = 1$. The Scherk-Kemperman Theorem is a basic tool in the proof by Gao that a sequence of elements of a finite abelian group G with length $|G| + d(G)$ contains a $|G|$ -sub-sequence summing to 0, where $d(G)$ is the maximal size of a sequence of elements of G having no non-empty zero-sum subsequence [2]. A recent generalization of Gao's Theorem, based also on the Scherk-Kemperman Theorem, is contained in [8]. Let G be a group and let B be a finite subset with $1 \notin B$. Using the Scherk-Kemperman Theorem, Eliahou and Lecouvey proved in [1] that there is a permutation σ of B such that $x\sigma(x) \notin B$, for every $x \in B$.

By a *graph* we shall mean a relation. Let $\Gamma = (V, E)$ be a graph and let F be a subset of V . As a consequence of this definition of a graph, $\Gamma(F)$ is just the image of F by the relation Γ , defined in elementary Set Theory. The graph $\Gamma = (V, E)$ is said to be *locally-finite* if $\Gamma(x)$ is finite for all $x \in V$.

Suppose now that Γ is a loopless finite graph with a transitive group of automorphisms. Let $v \in V$ and put $r = |\Gamma(v)|$. Motivated by some conjectures from Graph Theory, Mader [10], proved that there are directed cycles $C_1, \dots, C_r$ such that $C_i \cap C_j = \{v\}$, for all $i < j$.

After reading the present work, the reader could certainly use Menger's Theorem to prove that Mader's Theorem applied to Cayley graphs, is equivalent to the Scherk-Kemperman Theorem restricted to finite groups.

Notice that Mader's formulation fails in the infinite case, since infinite graphs with a transitive group of automorphisms could be acyclic. As a main tool, Mader introduced the notion of a vertex-fragment [10], mentioning some related difficulties.

Motivated by Moser's problem and by Mader's vertex-fragments, we shall define vertex-molecules. Our approach works in the infinite case too and leads to easier proofs. In particular, our approach avoids a duality between positive and negative vertex-fragments, in the spirit of the one introduced

in [4], used extensively in the arguments of Mader [10].

Let $\Gamma = (V, E)$ be a vertex-transitive reflexive locally-finite graph and let $v \in V$. Let F be a subset of V with $v \in F$.

Our main result states that

$$|\Gamma(F)| \geq |F| + |\Gamma(v)| - |\Gamma^-(v) \cap F|.$$

Applied to Cayley graphs, this result implies Kemperman's Theorem and the Scherk-Kemperman Theorem. We give also a simple proof of Mader's Theorem. Our result applies also to problems considered by Nathanson in [11] and by the author [5, 7].

2 Some Terminology

Let $\Gamma = (V, E)$ be a graph. The elements of V will be called *vertices*.

The *sub-graph* induced on a subset X is defined as $\Gamma[X] = (X, (X \times X) \cap E)$. The *degree* of a vertex x is defined as $d_\Gamma(x) = |\Gamma(x)|$. Our degree is called outdegree in some Graph Theory text books. Recall that the graph Γ is *locally-finite*, if Γ has only finite degrees. The graph Γ will be called *regular*, if all the vertices have the same degree.

The *boundary* of a subset X is defined as

$$\partial_\Gamma(X) = \Gamma(X) \setminus X.$$

We also write

$$\nabla_\Gamma(X) = V \setminus (X \cup \Gamma(X)).$$

When the context is clear, the reference to Γ will be omitted.

For any subset F of V , $V = F \cup \nabla(F) \cup \partial(F)$ is a partition (with possibly empty parts). Since $\Gamma(F) \cap \nabla(F) = \emptyset$, we have $F \cap \Gamma^-(\nabla(F)) = \emptyset$. Hence

$$\partial^-(\nabla(F)) \subset \partial(F). \tag{1}$$

The next lemma contains a useful sub-modular inequality:

Lemma A [6] *Let $\Gamma = (V, E)$ be a locally-finite reflexive graph and let X, Y be finite subsets of V . Then*

$$|\partial(X \cup Y)| + |\partial(X \cap Y)| \leq |\partial(X)| + |\partial(Y)|. \quad (2)$$

Proof. Observe that

$$\begin{aligned} |\Gamma(X \cup Y)| &= |\Gamma(X) \cup \Gamma(Y)| \\ &= |\Gamma(X)| + |\Gamma(Y)| - |\Gamma(X) \cap \Gamma(Y)| \\ &\leq |\Gamma(X)| + |\Gamma(Y)| - |\Gamma(X \cap Y)| \end{aligned}$$

By subtracting the equation $|X \cup Y| = |X| + |Y| - |X \cap Y|$, we obtain (2). ■

A map $f : V \rightarrow V$ will be called a *homomorphism* if $\Gamma(f(x)) = f(\Gamma(x))$, for every $x \in V$. A bijective homomorphism is called an *automorphism*. The graph Γ will be called *vertex-transitive* if for all $x, y \in V$, there is an automorphism f such that $y = f(x)$. Clearly a vertex-transitive graph is regular.

Let G be group and let F and S be subsets of G . The *Cayley graph* defined on G by S , is defined as $\text{Cay}(G, S) = (G, E)$, where $E = \{(x, y) : x^{-1}y \in S\}$. Notice that left translations are automorphisms of $\text{Cay}(G, S)$. In particular, Cayley graphs are vertex-transitive.

Putting $\Gamma = \text{Cay}(G, S)$, we have clearly $\Gamma(F) = FS$. Notice that Moser's condition " $A \cap B^{-1} = \{1\}$ " is just " $A \cap \Gamma^{-}(1) = \{1\}$ ".

3 Moser's sets

We investigate in this section a generalization of Moser's problem to graphs.

Let $\Gamma = (V, E)$ be a locally-finite reflexive graph and let $v \in V$. A set F is said to be a v -Moser's set if $\Gamma^{-}(v) \cap F = \{v\}$. Put

$$\mu(v, \Gamma) = \min\{|\partial(X)| : X \text{ is a } v\text{-Moser's set of } \Gamma\}.$$

A Moser's set X with $|\partial(X)| = \mu(v, \Gamma)$ will be called a v -*molecule*. The reference to Γ could be implicit.

Lemma 1 *Let $\Gamma = (V, E)$ be a locally-finite reflexive graph and let $v \in V$. Then the intersection and the union of two v -molecules are v -molecules.*

Proof.

Notice that the intersection and the union of two v -Moser sets are v -Moser sets. Let F_1 and F_2 be two v -molecules. Using (2), we have

$$\begin{aligned} 2\mu(v) &\leq |\partial(F_1 \cap F_2)| + |\partial(F_1 \cup F_2)| \\ &\leq |\partial(F_1)| + |\partial(F_2)| \leq 2\mu(v). \end{aligned}$$

Thus, $F_1 \cap F_2$ and $F_1 \cup F_2$ are v -molecules. ■

In particular, there exists a v -molecule contained in every v -molecule. Such a v -molecule will be called the v -kernel and denoted by K_v .

In the finite case, one may prove that a Mader's v -fragment [10] is either a v -molecule or a set obtained by deleting v from a v -molecule. Also the Mader's v -atom [10] is just $A_v = K_v \setminus \{v\}$. However our approach leads to simplifications, since we do not need negative v -fragments used by Mader. Notice that negative v -fragments have no nice behavior in the infinite case.

Lemma 2 *Let $\Gamma = (V, E)$ be a locally-finite reflexive graph and let $v, w \in V$ be vertices. If $v \neq w$, then $K_v \neq K_w$.*

Moreover $\phi(K_v) = K_{\phi(v)}$, for any automorphism ϕ of Γ .

Proof. Assume that $v \neq w$ and that $K_v = K_w$. It follows that $\Gamma^-(w) \cap K_v = \{w\}$. Thus $w \notin \partial(K_v \setminus \{w\})$. Using the definition of μ , we have

$$\mu(v) \leq |\partial(K_v \setminus \{w\})| \leq |\partial(K_v)| = \mu(v).$$

It would follow that $K_v \setminus \{w\}$ is a v -molecule, a contradiction.

Clearly $\phi(K_v)$ is a $\phi(v)$ -Moser's set. In particular, $K_{\phi(v)} \subset \phi(K_v)$. The reverse inclusion follows since $\phi^{-1}(K_{\phi(v)})$ is a v -Moser's set. ■

The next Lemma generalizes to the infinite case a lemma of Mader [10]. Note that Mader's argument is not suitable in the infinite case, since it involves negative v -fragments.

Lemma 3 *Let $\Gamma = (V, E)$ be a locally-finite reflexive graph and let $v, w \in V$ be vertices. If $w \in K_v$, then either $v \in \Gamma(K_w)$ or $K_w \subset K_v$.*

Proof. Suppose that $w \in K_v$ and that $v \in \nabla(K_w)$. We have using (1),

$$\begin{aligned}\Gamma^-(v) \cap \Gamma^-(\nabla(K_w)) &= \nabla(K_w) \cup \partial^-(\nabla(K_w)) \\ &\subset \nabla(K_w) \cup \partial(K_w) = V \setminus K_w.\end{aligned}$$

Therefore $\Gamma^-(v) \cap (K_v \cup K_w) = \Gamma^-(v) \cap K_v = \{v\}$. Thus $K_v \cup K_w$ is a v -Moser's set. Notice that $K_v \cap K_w$ is a w -Moser's set.

By (2),

$$\begin{aligned}\mu(v) + \mu(w) &\leq |\partial(K_v \cup K_w)| + |\partial(K_v \cap K_w)| \\ &\leq |\partial(K_v)| + |\partial(K_w)| \leq \mu(v) + \mu(w).\end{aligned}$$

Thus, $K_v \cap K_w$ is a w -molecule and hence $K_w \subset K_v$, a contradiction. ■

The *kernel-graph* Ω_Γ , introduced in the finite case by Mader in [10], is a graph on V with

$$\Omega_\Gamma(v) = \Gamma(v) \cap K_v,$$

for every $v \in V$. The reference to Γ could be implicit. The following easy lemma generalizes a result, proved by Mader [10] in the finite case:

Lemma 4 *Let $\Gamma = (V, E)$ be a vertex-transitive reflexive locally-finite graph. Then the kernel-graph Ω_Γ is a vertex-transitive graph. Moreover*

$$\Omega^-(v) \setminus \{v\} \subset \partial(K_v), \tag{3}$$

for every $v \in K_v$.

Proof. Take an automorphism ϕ of Γ . By Lemma 2, we have,

$$\phi(\Omega(v)) = \phi(\Gamma(v) \cap K_v) = \Gamma(\phi(v)) \cap K_{\phi(v)} = \Omega(\phi(v)),$$

showing that ϕ is an automorphism of Ω . Thus, Ω is vertex-transitive.

Take $w \in \Omega^-(v)$, with $w \neq v$. Take an automorphism ϕ with $\phi(v) = w$. We have by Lemma 2, $\phi(K_v) = K_w$, and hence $|K_v| = |K_w|$. By Lemma 2, $K_v \not\subset K_w$. By Lemma 3, $v \in \Gamma(K_w)$. Since $v \in \Gamma(w)$, we have $w \notin K_v$. ■

4 A first step

We prove here a special case of the main result. This special case implies the Scherk-Kemperman Theorem and Mader's Theorem.

Theorem 5 *Let $\Gamma = (V, E)$ be a vertex-transitive reflexive locally-finite graph and let $v \in V$. Then $\mu(v) \geq |\Gamma(v)| - |\Omega(v)| + |\Omega^-(v)| - 1$.*

Proof. Take an $x \in \Omega^-(v) \setminus \{v\}$. We have $v \in K_x$. By the definition of a Moser's set, $x \notin \Gamma(v)$. Thus,

$$\Omega^-(v) \cap \Gamma(v) = \{v\}.$$

By (3),

$$\Omega^-(v) \setminus \{v\} \subset \partial(K_v).$$

Clearly $\Gamma(v) \setminus K_v \subset \partial(K_v)$.

It follows that

$$\mu(v) = |\partial(K_v)| \geq |\Gamma(v) \setminus K_v| + |\Omega^-(v)| - 1 = |\Gamma(v)| - |\Omega(v)| + |\Omega^-(v)| - 1.$$

■

Let $\Gamma = (V, E)$ be a vertex-transitive locally-finite graph and let $v \in V$. If V is finite or if Γ is a Cayley graph, then $|\Gamma(v)| = |\Gamma^-(v)|$. The last well known fact is an easy exercise. In order to deduce the Scherk-Kemperman Theorem, we need to show that the kernel graph of a Cayley graph is also a Cayley graph.

Corollary 6 *(The Scherk-Kemperman Theorem [9]) Let A and B be a finite subsets of a group G with $A \cap (B^{-1}) = \{1\}$. Then $|AB| \geq |A| + |B| - 1$.*

Proof. Put $\Gamma = \text{Cay}(G, B)$. Notice that Γ is a reflexive vertex-transitive graph and that A is a 1-Moser's set of Γ .

We have $\Omega(x) = K_x \cap \Gamma(x) = xK_1 \cap xS = x(K_1 \cap S)$, by Lemma 2 and since a left translation is Cayley graph automorphism. Thus

$$\Omega_\Gamma = \text{Cay}(G, B \cap K_1).$$

It follows that

$$|AB| - |A| \geq \mu(1) \geq |\Gamma(1)| - |\Omega(1)| + |\Omega^-(1)| - 1 = |\Gamma(1)| - 1 = |B| - 1.$$

■

Corollary 7 *Let $\Gamma = (V, E)$ be a reflexive finite vertex-transitive graph and let $v \in V$. Let F be a subset of V with $v \in F$. Then*

$$|\Gamma(F)| \geq |F| + |\Gamma(v)| - |\Gamma^-(v) \cap F|.$$

Proof.

Assume first that $|\Gamma^-(v) \cap F| = 1$. Clearly F is a v -Moser's set.

By Lemma 4, Ω_Γ is a vertex-transitive graph. Since V is finite, we have $|\Omega(v)| = |\Omega^-(v)|$. By Theorem 5,

$$|\Gamma(F)| - |F| \geq \mu(v) \geq |\Gamma(v)| - |\Omega(v)| + |\Omega^-(v)| - 1 = |\Gamma(v)| - 1.$$

Assume now that $|\Gamma^-(v) \cap F| \geq 2$. Put $F' = (F \setminus (\Gamma^-(v))) \cup \{v\}$. By the first case, $|\Gamma(F)| \geq |\Gamma(F')| \geq |F'| + |\Gamma(v)| - 1 = |F| + |\Gamma(v)| - |\Gamma^-(v) \cap F|$. ■

For the next result, we assume some familiarity with Menger's Theorem and with the notion of a directed cycle in a graph.

Corollary 8 (Mader [10]) *Let $\Gamma = (V, E)$ be a vertex-transitive loopless finite graph and let $v \in V$. Put $|\Gamma(v)| = r$. Then there are directed cycles $C_1, \dots, C_r$ such that $C_i \cap C_j = \{v\}$, for all $i < j$.*

Proof. Let Φ be the graph obtained by adding vertex $v' \notin V$, with $\Phi(v') = \Gamma^-(v)$. Let Θ be the reflexive closure of Φ (obtained from Φ by adding loops everywhere). Let F be a subset of V with $v \in F$ and $v' \notin \Theta(F)$. Then clearly F is a v -Moser's set of Θ . By Theorem 10,

$$|\partial_\Phi(F)| = |\partial_\Theta(F)| \geq |\Theta(v)| - 1 = r.$$

By Menger's Theorem, Φ contains r disjoint paths from $\Gamma(v)$ to $\Gamma^-(v') = \Gamma^-(v)$. Adding v to each of these paths, we see the existence of r cycles with the desired property. ■

5 The main result

Let $\Gamma = (V, E)$ be a locally-finite reflexive graph. We define the *weak connectivity* of Γ as

$$\kappa_0(\Gamma) = \min\{|\partial(X)| : 1 \leq |X| < \infty\}. \quad (4)$$

This concept is not interesting in the finite case, since $\kappa_0(\Gamma) = 0$, for any finite graph Γ . A subset X achieving the minimum in (4) is called a *weak fragment* of Γ . A weak fragment with minimum cardinality will be called a *weak atom*.

Proposition 9 *Let $\Gamma = (V, E)$ be a locally-finite reflexive vertex-transitive graph and let A be a weak atom. Then the subgraph $\Gamma[A]$ induced on A is a vertex-transitive graph. Moreover every vertex belongs to some weak atom.*

Proof. Let F_1 and F_2 are weak fragments with $F_1 \cap F_2 \neq \emptyset$.

By (2), $2\kappa_0(\Gamma) \leq |\partial(F_1 \cap F_2)| + |\partial(F_1 \cup F_2)| \leq |\partial(F_1)| + |\partial(F_2)| \leq 2\kappa_0(\Gamma)$. Hence $F_1 \cap F_2$ is a weak fragment. In particular, distinct weak atoms are disjoint. Take $v \in A$. For every $y \in V$, there is an automorphism ϕ such that $\phi(v) = y$. Hence y belongs to the atom $\phi(A)$. If $y \in A$, then $\phi(A) = A$, and therefore ϕ/A is an automorphism of $\Gamma[A]$ with $\phi/A(v) = y$. Thus $\Gamma[A]$ is a vertex-transitive graph. ■

We are now ready to prove the general form of the main result:

Theorem 10 *Let $\Gamma = (V, E)$ be a reflexive locally-finite vertex-transitive graph and let $v \in V$. Let F be a finite subset of V with $v \in F$. Then*

$$|\Gamma(F)| \geq |F| + |\Gamma(v)| - |\Gamma^-(v) \cap F|.$$

Proof.

The result holds by Corollary 7, if V is finite. Suppose that V is infinite. By Proposition 9, there is a weak atom A with $v \in A$. By Proposition 9, $\Gamma[A]$ is a vertex-transitive graph. Clearly $F \cap A$ is Moser's set of the finite graph $\Gamma[A]$. By Corollary 7,

$$|\partial(F \cap A)| \geq |A \cap \Gamma(v)| - |A \cap \Gamma^-(v)|.$$

By the definition of κ_0 , we have $|\partial(F \cup A)| \geq \kappa_0 = |\partial(A)|$. Notice that $\Gamma(v) \subset A \cup \partial(A)$ and that $\partial(F \cup A) \setminus \partial(F) \subset \partial(A) \setminus \Gamma(v)$. It follows that

$$\begin{aligned}
|\partial(F)| &\geq |\partial(F \cap A) \cap A| + |\partial(F \cap A) \cap \partial(F)| \\
&\geq |\partial(F) \cap A| + |\partial(F \cup A)| - |\partial(A) \setminus \Gamma(v)| \\
&\geq |A \cap \Gamma(v)| - |A \cap \Gamma^-(v)| + |\partial(A)| - |\partial(A) \setminus \Gamma(v)| \\
&= |A \cap \Gamma(v)| - |A \cap \Gamma^-(v)| + |\partial(A) \cap \Gamma(v)| \geq |\Gamma(v)| - |F \cap \Gamma^-(v)|.
\end{aligned}$$

■

Corollary 11 (*Kemperman's Theorem [9]*)

Let A and B be finite subsets of a group G and let $c \in AB$. Then $|AB| \geq |A| + |B| - |A \cap (cB^{-1})|$.

Proof. Take $c = ab$, with $a \in A$ and $b \in B$. Put $A' = a^{-1}A$ and $B' = Bb^{-1}$. Put $\Gamma = \text{Cay}(G, B')$. Notice that Γ is a reflexive vertex-transitive graph and that A' is a 1-Moser's set of Γ . By Theorem 10,

$$\begin{aligned}
|AB| = |A'B'| &= |\Gamma(A')| \\
&\geq |A'| + |\Gamma(1)| - |A' \cap \Gamma^{-1}(1)| \\
&= |A'| + |B'| + |A' \cap B'^{-1}| = |A| + |B| - |A \cap (cB^{-1})|.
\end{aligned}$$

■

References

- [1] S. Eliahou and C. Lecouvey, Matchings in arbitrary groups, *Advances in Applied Mathematics* 40 (2008), 219-224.
- [2] W. D. Gao, A combinatorial problem on finite abelian groups. *J. Number Theory* 58 (1996), no. 1, 100-103.
- [3] A. Geroldinger, F. Halter-Koch, Non-unique factorizations. Algebraic, combinatorial and analytic theory. Pure and Applied Mathematics (Boca Raton), 278. Chapman & Hall/CRC, Boca Raton, FL, 2006. xxii+700 pp.
- [4] Y.O. Hamidoune, Sur les atomes d'un graphe orienté, C.R. Acad. Sc. Paris A 284 (1977), 1253-1256.

- [5] Y.O. Hamidoune, An application of connectivity theory in graphs to factorizations of elements in groups, *Europ. J of Combinatorics* 2 (1981), 349-355.
- [6] Y.O. Hamidoune, On small subset product in a group. *Structure Theory of set-addition*, *Astérisque*. no. 258(1999), xiv-xv, 281-308.
- [7] Y. O. Hamidoune. On iterated image size for point-symmetric relations. *Combin. Probab. Comput.* 17 (2008), no. 1, 61–66.
- [8] Y. O. Hamidoune, A weighted generalization of Gao's $n + D - 1$ -Theorem, *Combin. Probab. Comput.* 17 (2008), no. 17, 793–798.
- [9] J.H.B. Kemperman, On complexes in a semigroup, *Nederl. Akad. Wetensch. Proc. Ser. A.* 59 *Indag. Math.* 18 (1956), 247–254.
- [10] W. Mader, Existence of openly disjoint circuits through a vertex. *J of Graph Theory*, to appear.
- [11] M. B. Nathanson, The Caccetta-Häggkvist conjecture and Additive Number Theory, *arXiv:math. CO/0603469*, 2006.
- [12] J.E. Olson, Sums of sets of group elements, *Acta Arith.* 28 (1975/76), no. 2, 147–156.
- [13] P. Scherk, *Distinct elements in a set of sums*, *Amer. Math. Monthly*, 62 (1955), pp. 46–47.